

# Solitary Waves for the Generalized Nonautonomous Dual-power Nonlinear Schrödinger Equations with Variable Coefficients\*

Jin Gao<sup>1</sup>, Lijia Han<sup>2,†</sup> and Yehui Huang<sup>2</sup>

**Abstract** In this paper, we study the solitary waves for the generalized nonautonomous dual-power nonlinear Schrödinger equations (DPNLS) with variable coefficients, which could be used to describe the saturation of the nonlinear refractive index and the solitons in photovoltaic-photorefractive materials such as LiNbO<sub>3</sub>, as well as many nonlinear optics problems. We generalize an explicit similarity transformation, which maps generalized nonautonomous DPNLS equations into ordinary autonomous DPNLS. Moreover, solitary waves of two concrete equations with space-quadratic potential and optical super-lattice potential are investigated.

**Keywords** Solitary waves, dual-power law, nonlinear Schrödinger equation, variable coefficients.

**MSC(2010)** 35A09, 35C07, 35C08.

## 1. Introduction

Phenomena in nonlinear optics and the Bose-Einstein condensates (BECs) are often described by nonlinear Schrödinger equations (NLS) [4–6, 9–11, 14, 16–18, 20, 23, 28]. For example, the wave phenomena observed in fluid dynamics, plasma and elastic media and optical fibers, etc. When we want to understand the physical mechanism of phenomena, exact solutions for the nonlinear Schrödinger equations have to be explored. Moreover, various types of the NLS with non-Kerr nonlinearities, which contain Kerr law, power law, parabolic law, dual-power law as well as the logarithmic law, and other varying potentials were studied by many researchers in [1–3, 7, 11–13, 15, 19, 22, 24–27, 30].

In this paper, we consider solitary waves of the 1D generalized nonautonomous dual-power nonlinear Schrödinger equations with variable coefficients (DPNLS)

$$iQ_t + D(x, t)Q_{xx} + (lR_1(x, t)|Q|^n + kR_2(x, t)|Q|^{2n})Q + V(x, t)Q = 0, \quad (1.1)$$

where  $Q(x, t)$  is the complex envelope of the propagating beam of the modes,  $x$  is the propagation distance, and  $t$  is the retarded time,  $l, n, k$  are arbitrary constants,

<sup>†</sup>the corresponding author.

Email address: [gao-j17@mails.tsinghua.edu.cn](mailto:gao-j17@mails.tsinghua.edu.cn) (J. Gao), [hljmath@ncepu.edu.cn](mailto:hljmath@ncepu.edu.cn) (L. Han), [yhhuang@ncepu.edu.cn](mailto:yhhuang@ncepu.edu.cn) (Y. Huang)

<sup>1</sup>Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China

<sup>2</sup>Department of Mathematics and Physics, North China Electric Power University, Beijing 102206, China

\*Han was supported by the Fundamental Research Funds for the Central Universities (Grant 2018MS054).

$D(x, t)$  is the dispersion coefficient,  $R_1(x, t)$ ,  $R_2(x, t)$  are the dual-power nonlinearity coefficients, respectively, and  $V(x, t)$  is the external potential. This model contains many special types of the NLS with variable coefficients such as the cubic NLS equation, the cubic-quintic(CQ) NLS model, the generalized NLS model, etc [28]. Let  $n = 2, k = 0, l$  is still arbitrary constants, (1.1) collapses to the Kerr law nonlinear Schrödinger equations with variable coefficients

$$iQ_t + D(x, t)Q_{xx} + lR_1(x, t)|Q|^2Q + V(x, t)Q = 0, \quad (1.2)$$

which relates to Eq.(1) of [9]. Let  $l = 0, n, k$  is still arbitrary constants, (1.1) collapses to the power law nonlinear Schrödinger equations with variable coefficients

$$iQ_t + D(x, t)Q_{xx} + kR_2(x, t)|Q|^{2n}Q + V(x, t)Q = 0, \quad (1.3)$$

which relates to Eq.(1) of [19] and Eq.(2) of [24]. Let  $n = 2, l, k$  are still arbitrary constants, (1.1) collapses to the parabolic law or CQ law nonlinear Schrödinger equations with variable coefficients

$$iQ_t + D(x, t)Q_{xx} + (lR_1(x, t)|Q|^2 + kR_2(x, t)|Q|^4)Q + V(x, t)Q = 0, \quad (1.4)$$

which relates to Eq.(1) of [11] with  $\gamma(x, t) = 0$ . If  $V(x, t) = 0, D(x, t), R_1(x, t), R_2(x, t)$  are constant coefficients, (1.1) become an autonomous DPNLS equations, which describes the saturation of the nonlinear refractive index, and also serves as a basic model to describe the solitons in photovoltaic-photorefractive materials such as LiNbO3 [2]. However, if  $V(x, t), D(x, t), R_1(x, t), R_2(x, t)$  are the functions of  $x$  and  $t$  respectively, we can't obtain solutions by using the traveling wave method of [27]. To deal with this problem, we hope to use the similarity transformation to obtain solitary waves of (1.1).

In recent years, the similarity transformation has been used to obtain the solution of the nonlinear Schrödinger equations. In [9, 11], the authors solve the nonlinear Schrödinger equations with variable coefficients by using the similarity transformation. In [6–8], the authors also skillfully obtain solitary wave for the coupled nonlinear Schrödinger equations with variable coefficients by using the similarity transformation. In our paper, we firstly deduce the generalized nonautonomous DPNLS to the usual autonomous DPNLS equation (2.2) and obtain stationary solitary waves of the usual autonomous DPNLS (2.2), then using Galilean invariance, we can obtain moving solitary waves of (2.2). Finally using similarity transformation, we could obtain solitary waves of (1.1), so we only need to study solitary waves of (2.2).

Equation (1) can be viewed as the evolution equation  $Q_t = \frac{\delta \mathcal{H}}{\delta(iQ^*)}$ , where  $\mathcal{H}$  is the Hamiltonian function

$$\mathcal{H} = - \int_{-\infty}^{+\infty} [Q^* D(x, t)Q_{xx} + lR_1(x, t)|Q|^{n+2} + kR_2(x, t)|Q|^{2n+2} + V(x, t)|Q|^2] dx.$$

Generally speaking, the Hamiltonian  $\mathcal{H}$  in our model (1) is not conserved. It will be seen that this situation would be changed by employing a similarity transformation technique, and the Hamiltonian  $\mathcal{H}$  can only be conserved under some special cases.

The organization of this paper is as follows: In Section 2, we map the generalized nonautonomous DPNLS into the DPNLS by using the similarity transformation, then through some computation, we get stationary solitary waves of the

nonautonomous DPNLS. In Section 3, we design two interesting external potentials such as space-quadratic potential and optical super-lattice potential with DP nonlinearity, which have strong physical meanings in the nonlinear science. Moreover, we obtain solitary waves for these two interesting nonautonomous DPNLS with different values of  $k, l, n$ .

## 2. Solitary waves of nonautonomous NLSE

### 2.1. The similarity transformation

In this section, the generalized nonautonomous DPNLS (1.1) will be reduced into the autonomous DPNLS by the similarity transformation. In other words, solitary waves of the generalized nonautonomous DPNLS can be written as

$$Q(x, t) = q(X, T)p(x, t)e^{i\phi(x, t)}, \quad (2.1)$$

where  $X = X(x, t)$ ,  $T = T(t)$ ,  $q, p, \phi$  are real smooth functions of  $x$  and  $t$ . Using (2.1), we can reduce the generalized nonautonomous DPNLS (1.1) into the usual autonomous DPNLS

$$iq_T + q_{XX} + (l|q|^n + k|q|^{2n})q = 0, \quad (2.2)$$

where  $q(X, T)$  is solution of (2.2),  $l|q|^n + k|q|^{2n}$  is dual-power nonlinearity. In the following computation process, if  $T$  is only expressed by  $t$ , we can eliminate  $q_{XT}$  and obtain (2.2) by setting different function  $D, R_1, R_2, V$ . Take (2.1) into (1.1), we obtain

$$\begin{aligned} &ie^{i\phi}(pq_X X_t + pq_T T_t + p_t q + ipq\phi_t) + De^{i\phi}(2p_x q_X X_x + 2ipq_X X_x \phi_x + pq_{XX} X_x^2 \\ &+ pq_X X_{xx} + 2ip_x q\phi_x + p_{xx} q - pq\phi_x^2 + ipq\phi_{xx}) + (lR_1|pqe^{i\phi}|^n + kR_2|pqe^{i\phi}|^{2n})pqe^{i\phi} \\ &+ Vqe^{i\phi} = 0, \end{aligned} \quad (2.3)$$

where  $D = D(x, t)$ ,  $R_1 = R_1(x, t)$ ,  $R_2 = R_2(x, t)$ ,  $\phi = \phi(x, t)$  are arbitrary functions of  $x$  and  $t$ . In order to mapping (1.1) to (2.2), let

$$D = \frac{T_t}{2X_x^2}, \quad R_1 = \frac{T_t}{p^n}, \quad R_2 = \frac{T_t}{p^{2n}}, \quad (2.4)$$

then take (2.4) into (2.3), we obtain

$$\begin{aligned} &iq_T + q_{XX} + (l|q|^n + k|q|^{2n})q + \left(\frac{iq_X X_t}{T_t} + \frac{ip_t q}{pT_t} - \frac{q\phi_t}{T_t}\right) + \left(\frac{p_x q_X}{pX_x} + \frac{iq_X \phi_x}{X_x} + \frac{q_X X_{xx}}{2X_x^2}\right. \\ &\left. + \frac{ip_x q\phi_x}{pX_x^2} + \frac{p_{xx} q}{2pX_x^2} - \frac{q\phi_x^2}{2X_x^2} + \frac{iq\phi_{xx}}{2X_x^2}\right) + \frac{Vq}{T_t} = 0, \end{aligned} \quad (2.5)$$

or they also can be written as

$$iq_T + q_{XX} + (l|q|^n + k|q|^{2n})q + \frac{(ik_1 + k_2)q_X}{2pX_x^2 T_t} + \frac{(ik_3 + k_4)q}{2pX_x^2 T_t} = 0, \quad (2.6)$$

where  $k_i$ ,  $i = 1, 2, 3, 4$  are given by

$$k_1 = 2pX_x^2 X_t + 2pX_x \phi_x T_t, \quad (2.7)$$

$$k_2 = 2p_x X_x T_t + p X_{xx} T_t, \quad (2.8)$$

$$k_3 = 2p_t X_x^2 + 2p_x \phi_x T_t + p \phi_{xx} T_t, \quad (2.9)$$

$$k_4 = p_{xx} T_t - 2p X_x^2 \phi_t - p \phi_x^2 T_t + 2V p X_x^2. \quad (2.10)$$

(2.6) is very close to (2.2), so we need to do an assumption  $k_1 = k_2 = k_3 = k_4 = 0$ , which is vital step to our transformation. From  $k_2 = 0$ , we obtain  $p^2 X_x = f_1^2(t)$ . From  $k_1 = 0$ , we obtain  $\phi_x = \frac{-X_x X_t}{T_t}$ . From  $k_3 = 0$ , we obtain  $\frac{p^2}{X_x} = \frac{1}{F^2(x)}$ , where  $F(x)$  is integral function about  $x$ . Furthermore, from the assumption, we obtain the expression of  $p, X, \psi$

$$p = \sqrt{\frac{f_1(t)}{F(x)}}, \quad X = \int F(x) f_1(t) dx + f_2(t), \quad (2.11)$$

$$\phi = \int \frac{(\int F(x) f_{1t} dx + f_{3t}) F(x) f_1(t)}{T_t} dx + f_3(t), \quad (2.12)$$

where  $f_1(t), f_2(t), f_3(t)$  are integral function of  $t$ . Moreover, take the expression of  $X, p, T$  in (2.11)-(2.12) into (2.4), we obtain

$$D = \frac{T_t}{2F^2 f_1(t)^2} \quad R_1 = T_t \left( \frac{F}{f_1(t)} \right)^{\frac{n}{2}}, \quad R_2 = T_t \left( \frac{F}{f_1(t)} \right)^{\frac{n}{2}}. \quad (2.13)$$

Now the expression of the external potential  $V(x, t)$  is given by  $k_4 = 0$

$$V = \frac{8f_1^2(t)\phi_t F^4(x) + 4T_t \phi_x^2 F^2(x) - 3T_t F_x^2 + 2T_t F F_{xx}}{8F^4(x) f_1^2(t)}. \quad (2.14)$$

From the equation (2.11)-(2.12), we find that the external potential  $V(x, t)$  isn't arbitrary function but relates to  $X, p, T, \phi$  which are decided by  $F(x), f_1(t), f_2(t), f_3(t), T$ .

Thus similarity transformations (2.1) indeed map the generalized nonautonomous DPNLS to the nautonomous DPNLS systems, we only need to obtain solution of the nautonomous DPNLS equation (2.2) in the next section. Equations (2.11)-(2.12) and (2.14) can be seen as integrability condition on the generalized nonautonomous DPNLS (1.1). External potential (2.14) and DP nonlinearity (2.13) have different expression by choices of arbitrary functions  $F(x), f_1(t) (F(x) f_1(t) > 0), f_2(t), f_3(t), T(t)$ .

## 2.2. Solitary waves of the autonomous DPNLS

In this section, we get the stationary solitary waves of the autonomous DPNLS. Using Galilean invariance, we obtain moving solitary waves of (2.2). The form of the stationary solitary waves of (2.2) can be written as

$$q(X, T) = u(X) e^{i\mu T}, \quad (2.15)$$

where  $\mu$  is a positive propagation constant.  $u(X)$  obeys the nonlinear ordinary differential equation (ODE)

$$u_{XX} - \mu u + (lu^n + ku^{2n})u = 0, \quad (2.16)$$

where  $u(X)$  is a localized real function. Multiplying (2.16) by  $u_X$  and integrating once, after simple computation, we obtain the explicit expression of  $u$  as

$$u(X) = \left[ \frac{A}{B + \cos h(CX)} \right]^{\frac{1}{n}}, \quad (2.17)$$

$$A = \frac{(2+n)\mu B}{l}, \quad B = \text{sign}(l) \left[ 1 + \frac{(n+2)^2 k \mu}{(n+1)l^2} \right]^{-\frac{1}{2}}, \quad C = n\sqrt{\mu},$$

where  $(4\mu k + l^2) > 0$  is the condition of existence of solitary wave [27]. Moreover, we can obtain precisely the soliton solution of equation (1.1) from the soliton solution of (2.17) and the transformation (2.1) and (2.15). The soliton solution of equation (1.1) is

$$Q(x, t) = \left[ \frac{A}{B + \cos h(CX)} \right]^{\frac{1}{n}} p(x, t) e^{i(\mu T + \phi(x, t))}. \quad (2.18)$$

In order to better illustrate the property of dynamical evolution of solution (2.18) of (1.1), we set different functions  $F(x), f_1(t), f_2(t), f_3(t), T(t)$ , simultaneously  $T_1 = 0, T_2 = 0, T_3 = 0, c_0 = 1$  and then obtain expressions of  $D, R_1, R_2, V$ , which must satisfy our condition (2.11)-(2.12) and (2.14).

### 3. Designed external potentials

In this section, we will show two interesting nonautonomous DPNLS, where the potential functions  $V$  in equations (1.1) are specially chosen by setting different functions  $F(x), f_1(t), f_2(t), f_3(t), T(t)$ . In the following, we mainly introduce nonautonomous DPNLS with space-quadratic potential and optical super-lattice potential, simultaneously, the corresponding (DP) nonlinearity have significant expressions, which have important physical applicants in nonlinear optics and BEC problems [6, 8, 9, 11].

#### 3.1. Space-quadratic potential

Firstly, from equations (2.11)-(2.14), we set  $F = F_0, f_1 = a_0 \text{sech}(t), f_2 = 1, f_3 = 1$  where  $F_0 a_0 > 0$ . Then, the  $X, p, \phi, T$  are designed as

$$X = F_0 a_0 \text{sech}(t)x + 1, \quad T = a_1 \tanh(t), \quad (3.1)$$

$$p = \sqrt{\frac{a_0}{F_0} \text{sech}(t)}, \quad \phi = -\frac{a_0^2 F_0^2}{2a_1} \tanh(t)x^2 + 1, \quad (3.2)$$


where  $a_1$  is constant. From (3.1)-(3.2), we obtain nonautonomous DPNLS

$$iQ_t + D(t)Q_{xx} + (lR_1(t)|Q|^n + kR_2(t)|Q|^{2n})Q + V(x, t)Q = 0, \quad (3.3)$$

where

$$D(t) = \frac{a_1}{F_0^2 a_0^2}, \quad R_1(t) = a_1 \text{sech}^2(t) \left( \frac{a_0}{F_0 \cosh(t)} \right)^{-\frac{n}{2}}, \quad (3.4)$$

$$R_2(t) = a_1 \text{sech}(t)^2 \left( \frac{a_0}{F_0 \cosh(t)} \right)^{-n}, \quad V(x, t) = F_0^2 a_0^3 \frac{2 \text{sech}^2(t) - 1}{2a_1} x^2, \quad (3.5)$$



**Figure 1.** Solitary waves of the nonautonomous DPNLS.(a), Space-quadratic potential. (b), solitary waves (3.6) with  $l = 1, k = 0, n = 2$  and  $a_0 = a_1 = F_0 = 1$ . (c), solitary waves (3.6) with  $l = -1, k = 1, n = 2$  and  $a_0 = a_1 = F_0 = 1$ . (d), solitary waves (3.6) with  $l = -1, k = 1, n = \frac{3}{2}$  and  $a_0 = a_1 = F_0 = 1$ .

where  $D, R_1, R_2$  are functions only of  $t$ .  $V$  is space-quadratic potential which is plotted in Fig.1–(a). The space-quadratic potential has been studied in [11] and is significant in the (BECs) and nonlinear optics [3]. Equation (1.1) is the generalized dual-power nonlinear Schrödinger equations DPNLS with time coefficients. Solitary waves (2.1) can be written as


$$Q(x, t) = \frac{A\sqrt{a_0 \operatorname{sech}(t)}}{\sqrt{F_0}(B + \cosh(C(F_0 a_0 \operatorname{sech}(t)x + 1)))} e^{i(\mu a_1 \tanh(t) - \frac{a_0^2 F_0^2}{2a_1} \tanh(t)x^2 + 1)}. \tag{3.6}$$

Profiles of  $|Q|^2$  are plotted in Fig.1(b)-(d) with different  $l, n, k$ . Meanwhile, from (3.5), the amplitude of solitary waves is  $\frac{A\sqrt{a_0 \operatorname{sech}(t)}}{\sqrt{F_0}(B+1)}$ , depending on the changes of the parameter  $f_1(t)$ . Therefore, we obtain solitary wave by choosing different  $l, n, k, a_0, a_1$ .

### 3.2. Optical super-lattice potential

Firstly, from equations (2.11)-(2.14), we set  $F = \frac{1}{b_0 \cos \omega_0 x + 1}$ ,  $f_1 = 1 + b_1 \cos \omega_1 t$ ,  $f_2 = t$ ,  $f_3 = t$ , where  $b_0 > 1$ ,  $b_1 > 0$ ,  $\omega_i \in \mathbb{R}$ ,  $i = 0, 1$ . Then,  $p, X, \phi, T$  are designed as

$$X = \frac{2b_1 \cos(\omega_1 t) + 2}{\omega_0 \sqrt{b_0^2 - 1}} \operatorname{arctanh}\left(\frac{b_0 - 1}{\sqrt{b_0^2 - 1}} \tan\left(\frac{\omega_0}{2} x\right)\right) + t, \quad T = t, \tag{3.7}$$



**Figure 2.** Solitary waves of the nonautonomous DP-NLS. (a), Fourier-synthesized lattice potential.(b), solitary waves (3.14) with  $l = 1, k = 0, n = 2$  and  $b_0 = b_1 = \frac{1}{2}, \omega_0 = \omega_1 = 1$ . (c), solitary waves (3.14) with  $l = -1, k = 1, n = 2$  and  $b_0 = b_1 = \frac{1}{2}, \omega_0 = \omega_1 = 1$ . (d), solitary waves (3.14) with  $l = -1, k = 1, n = \frac{3}{2}$  and  $b_0 = b_1 = \frac{1}{2}, \omega_0 = \omega_1 = 1$ .

$$p = \sqrt{(1 + b_1 \cos(\omega_1 t))(b_0 \cos(\omega_0 x) + 1)}, \quad (3.8)$$

$$\begin{aligned} \phi = & \frac{1}{\omega_0 \sqrt{b_0^2 - 1}} \int \left[ \frac{1}{b_0 \cos(\omega_0 x) + 1} (-2b_1 \omega_1 \sin(\omega_1 t) \operatorname{arctanh}\left(\frac{b_0 - 1}{\sqrt{b_0^2 - 1}} \tan\left(\frac{\omega_0}{2} x\right)\right) \right. \\ & \left. + \omega_0 \sqrt{b_0^2 - 1} (\cos(\omega_1 t) b_1 + 1) \right] dx + t. \end{aligned} \quad (3.9)$$

The variable coefficients in (1.1) are designed as

$$D = \left( \frac{b_0 \cos(\omega_0 x) + 1}{b_1 \cos(\omega_1 t) + 1} \right)^2, \quad (3.10)$$

$$R_1 = (1 + b_1 \cos(\omega_1 t))(b_0 \cos(\omega_0 x) + 1)^{-\frac{n}{2}}, \quad (3.11)$$

$$R_2 = (1 + b_1 \cos(\omega_1 t))(b_0 \cos(\omega_0 x) + 1)^{-n}, \quad (3.12)$$

$$V = \phi_t + \frac{1}{2} \phi_x^2 - \frac{b_0^2 \omega_0^2 \sin^2(\omega_0)}{8(1 + b_1 \cos \omega_1 t)^2} + \frac{2b_0 \omega_0^2 \cos(\omega_0 x)}{8(1 + b_1 \cos \omega_1 t)}, \quad (3.13)$$

where  $D, R_1, R_2, V$  are functions about space  $x$  and time  $t$ ,  $V$  is fourier-synthesized lattice potential which is plotted in Fig.2-(a). The Fourier-synthesized lattice potential resembles a periodic sequence of hills in the transverse and longitudinal directions. In [10, 11, 21, 29], the authors consider the Fourier-synthesized lattice potential which have important physics meanings. In our paper, we consider the

more widely Fourier-synthesized lattice potential. (1.1) with the Fourier-synthesized lattice potential (3.13) display nonuniform distribution both transverse and longitudinal directions in the nonlinear science.

Solitary waves (2.1) can be written as

$$Q(x, t) = \frac{A\sqrt{(1+b_1\cos(\omega_1 t))(b_0\cos(\omega_0 x)+1)}}{B + \cosh(C(2\frac{b_1\cos(\omega_1 t)+1}{\omega_0\sqrt{b_0^2-1}} \operatorname{arctanh}(\sqrt{\frac{b_0-1}{b_0+1}} \tan(\frac{\omega_0}{2}x)) + t))} e^{i(\mu t + \phi)}. \quad (3.14)$$

Profiles of  $|Q|^2$  are plotted in Fig.2(b)-(d) with different  $l, n, k$ . Meanwhile, from (3.2), the amplitude of solitary waves is  $\frac{A\sqrt{(1+b_1\cos(\omega_1 t))(b_0\cos(\omega_0 x)+1)}}{B+1}$ , depending on the parameters  $f_1(t), F(x)$ . Therefore, the amplitude will be affected by all the parameters  $l, n, k, b_0, b_1, \omega_0, \omega_2$ , which leads to the complex changes of the solitary wave.

## 4. Conclusion

In this paper, we analytically obtain the solitary waves for the generalized dual-power nonlinear Schrödinger equations (DPNLS) with variable coefficients by using the similarity transformation. We design two interesting external potential and DP nonlinearity, and obtain solitary waves of corresponding nonautonomous DPNLS, which have important physical meanings in the nonlinear science. The method we used in this paper can be extended to obtain solitary waves of the higher-dimensional generalized DPNLS and study their interaction properties. Stability of solitary waves for (1.1) with time coefficients will be further discussed in a separate paper. In addition, Based on some novel physical phenomena in nonlinear optics and the BECs, we predict that new exact solutions of the generalized DPNLS will exist, such as rogue waves.

## Acknowledgments

The authors would like to extend their sincere gratitude to Professor Liming Ling for their great help and advice during the research.

## References

- [1] A. Biswas, *Quasi-stationary non-Kerr law optical solitons*, Optical Fiber Technology, 2003, 9(4), 224–259.
- [2] A. Biswas and S. Konar, *Introduction to Non-Kerr Law Optical Solitons*, CRC Press, Boca Raton, FL, USA, 2007.
- [3] K. Bulent and I. Mustafa, *On optical solitons of the resonant Schrödinger's equation in optical fibers with dual-power law nonlinearity and time-dependent coefficients*, Waves in Random Media, 2015, 25(3), 8.
- [4] F. Dalfovo, S. Giorgini, L. Pitaevskii and S. Stringari, *Theory of trapped Bose-Condensed gases*, Reviews of Modern Physics, 1999, 71(3), 463.



- [5] B. Guo, L. Ling and Q. Liu, *High-order solutions and generalized darboux transformations of derivative nonlinear Schrödinger Equations*, Studies in Applied Mathematics, 2013, 130(4), 317–344.
- [6] B. Guo, L. Ling and Rogue Wave, *Breathers and Bright-Dark-Rogue solutions for the coupled Schrödinger equations*, Chinese Physics Letters, 2011, 28(11), 110202–110205.
- [7] L. Han and J. Gao, *Stability of solitary waves for the generalized nonautonomous dual-power nonlinear Schrödinger equations with time-dependent coefficients*, Communications in Nonlinear Science and Numerical Simulation, 2017, 42, 520–531.
- [8] L. Han, Y. Huang and H. Liu, *Solitons in coupled nonlinear Schrödinger equations with variable coefficients*, Communications in Nonlinear Science and Numerical Simulation, 2014, 19(9), 3063–3073.
- [9] J. He and Y. Li, *Designable integrability of the variable coefficient nonlinear Schrödinger equation*, Studies in applied mathematics, 2011, 126, 1–15.
- [10] J. He and H. Li, *Nonautonomous bright matter-wave solitons and soliton collisions in Fourier-synthesized optical lattices*, Optics Communications, 2011, 284(12), 3084–3089.
- [11] J. He and H. Li, *Analytical solitary-wave solutions of the generalized nonautonomous cubic-quintic nonlinear Schrödinger equation with different external potentials*, Physical Review E, 2011, 83(6), 066607.
- [12] L. Han and L. Xin, *Stability and soliton solutions for a parity-time-symmetric vector nonlinear Schrödinger system*, Indian Journal of Physics, 2018, 92, 1291–1298.
- [13] J. Jasinski, *New exact and approximate stationary solutions of (1+2)D nonlinear Schrödinger equation in dual-power media*, Optics Communications, 2015, 343, 80–84.
- [14] Y. Kivshar and G. Agrawal, *Optical Solitons: From Fibers to Photonic Crystals*, Academic Press, New York, 2003.
- [15] R. Kohl, A. Biswas, D. Milovic and E. Zerrad, *Optical soliton perturbation in a non-Kerr law media*, Optics and Laser Technology, 2008, 40(4), 647–662.
- [16] L. Ling and L. Zhao, *Simple determinant representation for rogue waves of the nonlinear Schrödinger equation*, Physical Review E, 2013, 88(4), 043201.
- [17] H. Luo, D. Zhao and X. He, *Exactly controllable transmission of nonautonomous optical solitons*, Physical Review A, 2009, 79(6), 1989–1990.
- [18] B. Malomed, D. Mihalache, F. Wise and L. Torner, *Spatiotemporal optical solitons*, Journal of Optics B Quantum and Semiclassical Optics, 2005, 7(5), R53.
- [19] R. Popovych, N. Ivanova and H. Eshraghi, *Group classification of (1+1)-dimensional Schrödinger equations with potentials and power nonlinearities*, Journal of Mathematical Physics, 2004, 45(8), 3049–3057.
- [20] C. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases*, Cambridge University Press, Cambridge, 2002.

- [21] T. Salger, C. Geckeler, S. Kling and M. Weitz, *Atomic Landau-Zener tunneling in Fourier-Synthesized optical lattices*, Physical Review Letters, 2007, 99(19), 190405.
- [22] V. Serkin, A. Hasegawa and T. Belyaeva, *Nonautonomous solitons in external potentials*, Physical Review Letters, 2007, 98(7), 074102.
- [23] C. Sulem and P. Sulem, *The Nonlinear Schrödinger Equation: Self-Focusing and Wave Collapse*, Springer-Verlag, Berlin, 1999.
- [24] A. Wazwaz, *Reliable analysis for nonlinear Schrödinger equations with a cubic nonlinearity and a power law nonlinearity*, Mathematical and Computer Modelling, 2006, 43(1), 178–184.
- [25] L. Wu, J. Zhang and L. Li, *Modulational instability and bright solitary wave solution for Bose-Einstein with time-dependent scattering length and harmonic potential*, New Journal of Physics, 2007, 9(3), 69.
- [26] L. Xin, Y. Kong and L. Han, *Solutions of solitary-wave for the variable-coefficient nonlinear Schrödinger equation with two power-law nonlinear terms*, International Journal of Modern Physics B. 2018, 32(28), 1850310.
- [27] J. Yang, *Nonlinear Waves in Integrable and Nonintegrable Systems*, SIAM, Philadelphia, 2010.
- [28] Z. Yan and V. Konotop, *Exact solutions to three-dimensional generalized nonlinear Schrödinger equations with varying potential and nonlinearities*, Physical Review E, 2009, 80(3), 036607.
- [29] Z. Yang, L. Zhao, T. Zhang, Y. Li and R. Yue, *Snakelike nonautonomous solitons in a graded-index grating waveguide*, Physical Review A, 2010, 81(4), 82–82.
- [30] X. Zheng and Y. Shang, *Abundant explicit exact solutions to the generalized nonlinear Schrödinger equation with parabolic law and dual-power law nonlinearities*, Mathematical Methods in the Applied Sciences, 2015, 38(2).